

Changes in Stratigraphic Nomenclature by the U.S. Geological Survey, 1971

By GEORGE V. COHEE *and* WILNA B. WRIGHT

CONTRIBUTIONS TO STRATIGRAPHY

GEOLOGICAL SURVEY BULLETIN 1372-A



CONTRIBUTIONS TO STRATIGRAPHY

CHANGES IN STRATIGRAPHIC NOMENCLATURE BY THE U.S. GEOLOGICAL SURVEY, 1971

By GEORGE V. COHEE and WILNA B. WRIGHT

LISTING OF NOMENCLATURAL CHANGES

In the following table, stratigraphic names adopted, revised, reinstated, or abandoned are listed alphabetically. The age of the unit, the revision, and the area involved, along with the author's name and date of publication of the report, are given. The publication in which the changes in nomenclature were made are listed in the references at the end of this publication. The capitalization of age terms in the age column follows official usage.

The following formal designations of Precambrian time are now in use by the U.S. Geological Survey:

Precambrian Z—base of Cambrian to 800 m.y.

Precambrian Y—800 m.y. to 1,600 m.y.

Precambrian X—1,600 m.y. to 2,500 m.y.

Precambrian W—older than 2,500 m.y.

On the recommendation of a Special Panel—consisting of Precambrian specialists M. D. Crittenden, Jr. (Chairman), J. E. Harrison, and J. C. Reed, Jr., working in collaboration with H. L. James—the Geologic Names Committee and the Chief Geologist approved an interim scheme for subdivision of Precambrian time, to be used in reports of the Geological Survey.

The basic control for selection of time boundaries and units is a correlation chart for the Precambrian of the United States, prepared by Z. E. Peterman and C. E. Hedge of the U.S. Geological Survey (unpub. data). The time boundaries have been chosen so as to split as few of the known epochs of sedimentation, orogeny, and plutonism as possible. The boundaries do not correspond intentionally to geologic events.

The scheme of subdivision has been devised simply to facilitate depiction and analysis of the Precambrian history of the United States.

The scheme is intended as an interim measure, pending development of an internationally accepted standard.

For depiction on maps, only the letter designations (W, X, Y, Z) will be shown as map symbols, and lowercase letters will indicate the group or formation names as appropriate. If a unit extends across the boundary between letter-designated units, both letters, the younger first, will be used in the map symbol. When geochronologic data are not adequate for unit assignment, only the general term Precambrian

Name	Age	Location
Aguas Buenas Limestone Member (of Fajardo Formation).	Early Cretaceous.....	Puerto Rico.....
Alder Group.....	Precambrian.....	Central Arizona.....
Almirante Sur Sand Member (of Cibao Formation).	late Oligocene and (or) early Miocene.	Puerto Rico.....
Alutom Formation.....	late Eocene, Oligocene, early Miocene.	Guam.....
Analomink Red Shale Member (of Catskill Formation).	Late Devonian.....	Northeastern Pennsylvania.
Angelica Arkose.....	Early Cretaceous.....	Southern Arizona....
Ardath Shale (of La Jolla Group).	middle Eocene.....	Southern California..
Ashdown Tuff.....	early Miocene.....	Northwestern Nevada.
Attalla Chert Conglomerate Member (of Chickamauga Limestone).	Middle Ordovician.....	Northwestern Georgia.
Baird Group.....	Middle and Late Devonian and older(?).	Northern Alaska.....
Bancroft Springs Basalt (of Snake River Group).	late Pleistocene.....	Southern Idaho.....
Bannock Volcanic Member (of Pocatello Formation).	late Precambrian.....	Southeastern Idaho (Pocatello area).
Barker Porphyry.....	Eocene(?).....	Montana.....
Barrazas Formation.....	Late Cretaceous (Cenomanian).	Northeastern Puerto Rico.
Bass Islands Dolomite.....	Late Silurian.....	Northern Ohio and southeastern Michigan.
Bates Mountain Tuff.....	Miocene.....	Nevada.....
Beaver Basin Formation.....	Pleistocene and Holocene	Utah.....
Bell Brook Formation.....	Silurian(?) and Devonian	Maine.....
Bena Gravel.....	Miocene.....	California.....
Big Blue Serpentinous Member (of Temblor Formation).	middle Miocene.....	California.....

and the symbol pC will be used. Rock units and events within a major time unit such as W, X, Y or Z, keyed to geochronologic data as available, will be shown on map explanations by simple sequential arrangement.

The previously used age designations for the Precambrian are given in the table because they were used by the authors in reports submitted to the Geologic Names Committee before the new scheme was adopted.

Revision and reference

Age changed from Late Cretaceous to Early Cretaceous. (Glover, 1971.)

Alder Group restricted to Mazatzal Mountains, its type locality, and removed from Yavapai Series. In Prescott-Jerome area, Big Bug Group (new) replaces Alder Group. (Anderson and others, 1971.)

Almirante Sur Sand Member changed to Almirante Sur Sand Lentil. (Monroe, 1971.)

Age changed from late Eocene and Oligocene to late Eocene, Oligocene, and early Miocene. (Bukry and others, 1971.)

Analomink Red Shale of Willard (1935) adopted as middle of three members of Catskill Formation. Overlies Delaware River Flags Member (not new); underlies Shohola Member (not new). (Alvord and Drake, 1971.)

Angelica Arkose adopted. Overlies Whitcomb Quartzite (new); underlies Demetrie Volcanics (not new). (Cooper, 1971.)

Ardath Shale adopted. Rocks were formerly part of Hanna's (1926) Rose Canyon Shale Member of La Jolla Formation of Clark (1926). Overlies Torrey Sandstone (not new); underlies Scripps Formation (new). (Kennedy and Moore, 1971.)

Ashdown Tuff adopted. (Noble and others, 1970.)

Geographically extended into northwestern Georgia and raised to formational rank as Attalla Conglomerate. (Cressler, 1970.)

Age changed from Middle(?) and Late Devonian to Middle and Late Devonian and older(?). (Brosge and Tailleux, 1971.)

Bancroft Springs Basalt abandoned; rocks now considered part of McKinney Basalt, of Snake River Group. (Malde, 1971.)

Bannock Volcanic Formation of Anderson (1928), as stratigraphically restricted by Ludlum (1942, 1943), adopted as Bannock Volcanic Member. Overlies unnamed lower member of Pocatello; underlies Scout Mountain Member (new). (Crittenden and others, 1971.)

Age changed from post-Cretaceous to Eocene(?). (Witkind, 1971.)

Barrazas Formation adopted; underlies Hato Puerco Formation. (Seiders, 1971.)

Bass Islands raised to group rank in Ohio and includes (ascending): Greenfield Dolomite, Tymochtee Formation, Put-in-Bay Dolomite, and Raisin River Dolomite. In Michigan, Bass Islands Dolomite (restricted) remains of formational rank and includes Put-in-Bay and Raisin River Dolomite Members. (Norris and Fidler, 1971.)

Age changed from Oligocene or Miocene to Miocene. (McKee and Stewart, 1971.)

Age changed from Pleistocene to Pleistocene and Holocene. (Birkeland and others, 1971.)

Age changed from Silurian(?) or Devonian(?) to Silurian(?) and Devonian. (Pavlidis, 1971.)

Bena Gravel of Dibblee and Chesterman (1953) adopted. (Dibblee and Warne, 1970.)

Name changed, removed from uppermost part of Temblor Formation, redefined by Adegoke (1969), and raised in rank to Big Blue Formation. Overlies Temblor Formation (now stratigraphically restricted) in New Idria area. (Lockwood, 1971.)

Name	Age	Location
Big Bug Group.....	Precambrian.....	Central Arizona.....
Blackhawk Formation (of Mesa-verde Group).	Late Cretaceous.....	Utah.....
Blackrock Canyon Limestone..	late Precambrian.....	Southeastern Idaho (Pocatello area).
Blue Hill Granite Porphyry....	Late Ordovician.....	Massachusetts.....
Borden Formation.....	Early and Late Mississippian.	Kentucky.....
Brigham Quartzite.....	late Precambrian to Middle Cambrian.	Northern Utah and southeastern Idaho.
Brodhead Creek Member (of Marcellus Shale).	Middle Devonian.....	Eastern Pennsylvania.
Bromley Shale Member (of Point Pleasant Formation).	Middle Ordovician.....	Northern Kentucky..
Browns Hole Formation.....	late Precambrian.....	Northern Utah (Huntsville area).
Buttermilk Falls Limestone....	Middle Devonian.....	Northeastern Pennsylvania.
Cabrillo Formation (of Rosario Group).	Late Cretaceous.....	Southern California..
Caddy Canyon Quartzite.....	late Precambrian.....	Southeastern Idaho (Pocatello area) and possibly northern Utah (Huntsville area).
Caetano Tuff.....	Oligocene.....	Nevada.....
Cambalache Formation.....	Late Cretaceous.....	Northeastern Puerto Rico.
Camelback Mountain Quartzite	late Precambrian and Early Cambrian.	Southeastern Idaho (Pocatello area).
Canóvanas Formation.....	Late Cretaceous.....	Northeastern Puerto Rico.
Cape Ann Granite.....	Late Ordovician.....	Massachusetts.....
Carter Sandstone Member (of Pierre Shale).	Late Cretaceous.....	North-central Colorado.
Carter Caves Sandstone.....	Late Mississippian.....	Northeastern Kentucky.
Catskill Formation.....	Late Devonian.....	Pennsylvania and New Jersey.
Celada Formation.....	Late Cretaceous.....	Northeastern Puerto Rico.
Chainman Shale.....	Late Mississippian.....	East-central Nevada..
Chainman Shale.....	Late Mississippian.....	East-central Nevada..
Chalk Mountain Nevadite.....	Eocene.....	Colorado.....
Chaparral Volcanics.....	Precambrian.....	Arizona.....

Revision and reference

- Big Bug Group adopted. Replaces Alder Group in Prescott-Jerome area. Includes (ascending): Green Gulch Volcanics, Spud Mountain Volcanics, and Iron King Volcanics. (Anderson and others, 1971.)
- Kenilworth and Sunnyside Members of Young (1955) of Blackhawk Formation adopted. Formation includes (ascending): Aberdeen, Kenilworth, lower mudstone, Sunnyside, and upper mudstone members. (Maberry, 1971.)
- Blackrock Canyon Limestone adopted. Same formation as Blackrock Limestone of Anderson (1928). Overlies Pocatello Formation (not new); underlies Papoose Creek Formation (new). (Crittenden and others, 1971.)
- Age changed from Devonian(?) to Late Ordovician. (Zartman and Marvin, 1971.)
- New Providence Shale reinstated in central Kentucky as basal member of Borden Formation. Nancy Member restricted to more silty, nonplastic shale within Nancy Member of previous usage. (Taylor and Lewis, 1971.)
- In its type area, Wasatch Mountains, Brigham is raised to group rank. Includes (ascending): Caddy Canyon(?) Quartzite (new), Inkorn(?) Formation (new), Mutual Formation, Browns Hole Formation (new), and Geertsen Canyon Quartzite (new). Brigham Quartzite remains in good usage in southeasternmost Idaho; Brigham restricted from usage in Pocatello area. (Crittenden and others, 1971.)
- Brodhead Creek Member of Willard (1938) adopted as upper of three members of Marcellus Shale. Overlies Stony Hollow Member (not new). (Alvord and Drake, 1971.)
- Bromley Shale of Bassler (1906) adopted as Bromley Shale Member of Point Pleasant Formation. (Luft, 1971.)
- Browns Hole Formation adopted; assigned to Brigham Group (newly revised). Overlies Mutual Formation; underlies Geertsen Canyon Quartzite (new). (Crittenden and others, 1971.)
- Buttermilk Falls Limestone of Willard (1939) adopted. Overlies Schoharie Formation; underlies Marcellus Shale. (Alvord and Drake, 1971.)
- Cabrillo Formation adopted as uppermost formation of Rosario Group (not new); overlies Point Loma Formation (new). (Kennedy and Moore, 1971.)
- Caddy Canyon Quartzite adopted. Overlies Papoose Creek Formation (new); underlies Inkorn Formation (new). In Huntsville area of northern Utah only, tentatively assigned as lowermost formation of Brigham Group (newly revised), overlying Kelley Canyon Formation (new) and underlying Inkorn(?) Formation of Brigham Group. (Crittenden and others, 1971.)
- Age changed from Oligocene(?) to Oligocene. (Wells and others, 1971.)
- Cambalache Formation adopted. Overlies Hato Puerco Formation; underlies Canóvanas Formation (new). Toma de Agua Vitrophyre Member (new) adopted with restricted occurrence at top of Cambalache. (Seiders, 1971.)
- Camelback Mountain Formation adopted. Overlies Mutual Formation; underlies Gibson Jack Formation (new). (Crittenden and others, 1971.)
- Canóvanas Formation adopted. Overlies Cambalache Formation (new); underlies Martín González Lava. (Seiders, 1971.)
- Age changed from late(?) Paleozoic to Late Ordovician. (Zartman and Marvin, 1971.)
- Carter Sandstone Member adopted; underlies and overlies two unnamed shale members of Pierre. (Izett and others, 1971.)
- Carter Caves Sandstone adopted as uppermost Mississippian formation (at Mississippian-Pennsylvanian systemic boundary) in northeastern Kentucky. (Englund and Windolph, 1971.)
- Delaware River Flags of White (1882) as used by Willard (1937), Anomink Red Shale of Willard (1935), and Shohola Formation of Willard (1936) adopted as basal, middle, and upper members, respectively, of Catskill Formation in report area. (Alvord and Drake, 1971.)
- Celada Formation adopted. Overlies Infierno Formation (new). (Seiders, 1971.)
- Age changed from Early and Late Mississippian to Late Mississippian in Diamond Peak area. (Gordon, 1971.)
- Name changed to Chainman Formation in southern Diamond Mountains. Chainman Shale remains in good usage elsewhere. (Brew, 1971.)
- Name changed to Chalk Mountain Rhyolite. (Bergendahl and Koschmann, 1971.)
- Chaparral Volcanics abandoned; its rocks now included in upper part of Spud Mountain Volcanics. (Anderson and others, 1971.)

Name	Age	Location
Chattanooga Shale.....	Devonian and Mississippian.	Northwestern Georgia.
Cheaha Sandstone Member (of Talladega Slate).	Paleozoic(?).....	Northwestern Georgia.
Chickamauga Limestone.....	Middle and Late Ordovician.	Northwestern Georgia.
Chirikof Formation.....	middle Tertiary.....	Southwestern Alaska.
Chopawamsic Formation.....	Cambrian or Ordovician.	North-central Virginia.
Chuniksak Formation.....	Miocene(?).....	Southwestern Alaska.
Church Creek Formation.....	Eocene (Refugian).....	California.....
Coldwater Shale.....	Early Mississippian.....	Michigan.....
Cornishville Bed (of Perryville Limestone Member) (of Lexington Limestone).	Middle Ordovician.....	Kentucky.....
Dakota Sandstone.....	Early(?) and Late Cretaceous.	New Mexico.....
Deaton Member (of Lenoir Limestone).	Middle Ordovician.....	Northwestern Georgia.
Delaware River Flags Member (of Catskill Formation).	Late Devonian.....	Northeastern Pennsylvania and southeastern New York.
Delmar Formation (of La Jolla Group).	middle Eocene.....	Southern California..
Demetrie Volcanics.....	Late Cretaceous.....	Southern Arizona....
Diamond Peak Formation.....	Early and Late Mississippian and Early Pennsylvanian.	East-central Nevada..
Draper Formation.....	Pleistocene and Holocene	Utah.....
Eagle Valley Evaporite.....	Middle Pennsylvanian..	Colorado.....
Echo Canyon Conglomerate....	Late Cretaceous.....	Northern Utah.....
Edwards Creek Tuff.....	Oligocene or Miocene...	Central Nevada.....
Elk Ridge Limestone Member (of Minturn Formation).	Middle Pennsylvanian..	Colorado.....
Ely Limestone.....	Late Mississippian and Early Pennsylvanian.	Nevada (Eureka area).
Everett Formation.....	Cambrian(?) and Cambrian.	Massachusetts, Connecticut, and New York.
Fajardo Formation.....	Early Cretaceous.....	Puerto Rico.....
Faneto Formation.....	late Tertiary or early Pleistocene.	Southwestern Alaska..
Farmers Member (of Borden Formation).	Early Mississippian.....	Eastern Kentucky....
Flowery Trail Granodiorite....	Mesozoic(?).....	Northeastern Washington.
Floyd Shale.....	Late Mississippian.....	Northwestern Georgia.
Fort Payne Chert.....	Early Mississippian.....	Northwestern Georgia.
Frailles Formation.....	Late Cretaceous.....	Northeastern Puerto Rico.

Revision and reference

Maury Formation reduced in rank to member of Chattanooga Shale in northwestern Georgia. (Cressler, 1970.)

Geographically extended into northwestern Georgia. (Cressler, 1970.)

Attalla Chert Conglomerate Member of Chickamauga Limestone raised to formation rank as Attalla Conglomerate and geographically extended into northwestern Georgia. (Cressler, 1970.)

Chirikof Formation adopted. (Gates and others, 1971.)

Chopawamsic Formation adopted. (Southwick and others, 1971.)

Chuniksak Formation adopted. (Gates and others, 1971.)

Church Creek Formation of Dickinson (1965) adopted. (Brabb and others, 1971.)

Age changed from Mississippian to Early Mississippian. (Sable, 1970.)

Cornishville Limestone Member (of Perryville Formation) of Foerste (1912) adopted as Cornishville Bed of Perryville Limestone Member of Lexington Limestone. (Wolcott and Cressman, 1971.)

Twowells Sandstone Lentil (Pike, 1947) of Mancos Shale adopted as Twowells Tongue of Dakota Sandstone. Whitewater Arroyo Shale Member of Owen (1966) of Dakota Sandstone adopted as Whitewater Arroyo Tongue of Mancos Shale. (Green and Pierson, 1971.)

Deaton Formation of Spencer (1893) adopted as Deaton Member of Lenoir Limestone. (Cressler, 1970.)

Delaware River Flags of White (1882), as used by Willard (1937), adopted as lower of three members of Catskill Formation. Underlies Analomink Red Shale Member (not new). (Alvord and Drake, 1971.)

Delmar Sand of Hanna (1926), member of La Jolla Formation of Clark (1926), adopted as Delmar Formation of La Jolla Group (not new). Overlies Mount Soledad Formation (new); underlies Torrey Sandstone (not new). (Kennedy and Moore, 1971.)

Demetrie Formation of Thoms (1967) adopted as Demetrie Volcanics. Truncates several Mesozoic formations; underlies Red Boy Rhyolite (not new). (Cooper, 1971.)

In northern Pinyon Range, age of Diamond Peak Formation is Early and Late Mississippian and Early Pennsylvanian; elsewhere, age is Late Mississippian. (Gordon, 1971.)

Age changed from Pleistocene to Pleistocene and Holocene. (Birkeland and others, 1971.)

Age changed from Pennsylvanian and Permian to Middle Pennsylvanian. (Mallory, 1971.)

Echo Canyon Conglomerate of Williams and Madsen (1959) adopted. (Mullens, 1971.)

Edwards Creek Tuff adopted. (McKee and Stewart, 1971.)

Age changed from Pennsylvanian to Middle Pennsylvanian. (Mallory, 1971.)

Age changed from Pennsylvanian to Late Mississippian and Early Pennsylvanian. (Nolan and others, 1971.)

Age changed from Cambrian(?), Cambrian, or Ordovician, to Cambrian(?) and Cambrian. (Zen and Ratcliffe, 1971.)

Age changed from Late Cretaceous to Early Cretaceous. (Glover, 1971.)

Faneto Formation adopted. (Gates and others, 1971.)

Age changed from Mississippian to Early Mississippian. (Sable, 1970.)

Flowers Trail Granodiorite adopted. (Clark and Miller, 1968.)

Hartselle Sandstone reduced in rank to member of Floyd Shale in northwestern Georgia. (Cressler, 1970.)

Lavender Shale Member of Butts and Gildersleeve (1948) of Fort Payne Chert adopted. (Cressler, 1970.)

Leprocomio Siltstone Member of Guaynabo Formation reassigned to Frailes Formation as Leprocomio Mudstone Member. (Seiders, 1971.)

Name	Age	Location
Friars Formation (of La Jolla Group).	middle and late Eocene..	Southern California..
Frog Mountain Sandstone.....	Early and Middle Devonian.	Northwestern Georgia.
Gasconade Dolomite.....	Early Ordovician.....	Missouri.....
Geertsen Canyon Quartzite....	late Precambrian and Early(?) Cambrian.	Northern Utah (Huntsville area).
Gibson Jack Formation.....	Early Cambrian.....	Southeastern Idaho (Pocatello area).
Golden Door Volcanics.....	Tertiary.....	Arizona and Nevada..
Goshen Formation.....	Middle and Late Silurian to Early Devonian.	Massachusetts.....
Granite Mountain Porphyry....	Paleocene.....	Southeastern Arizona.
Grant Mills Granodiorite.....	Mississippian(?) or older.	Rhode Island.....
Graydon Channel Sandstone....	Pennsylvanian.....	Missouri.....
Greenfield Dolomite Member (of Bass Islands Dolomite).	Late Silurian.....	Northern Ohio.....
Green Gulch Volcanics.....	Precambrian.....	Central Arizona.....
Guaracanal Andesite.....	Paleocene.....	Northeastern Puerto Rico.
Guaynabo Formation.....	Late Cretaceous.....	Northeastern Puerto Rico.
Gunsight Pass Member (of Pierre Shale).	Late Cretaceous.....	North-central Colorado.
Gunter Sandstone Member (of Van Buren Formation).	Early Ordovician.....	Missouri.....
Harkers Alluvium.....	Pleistocene.....	Utah.....
Hartselle Sandstone.....	Mississippian.....	Northwestern Georgia.
Hato Puerco Tuff.....	Late Cretaceous.....	Northeastern Puerto Rico.
Henderson Gneiss.....	late Precambrian or Paleozoic.	North Carolina.....
Henley Bed (of Farmers Member) (of Borden Formation).	Early Mississippian....	Eastern Kentucky...
Hickey Formation.....	late Miocene and Pliocene(?).	Central Arizona.....
Hickey Formation.....	late Miocene and early Pliocene.	Central and north-central Arizona.
Hillabee Chlorite Schist.....	post-Carboniferous....	Northwestern Georgia.
Hornsilver Dolomite Member (of Minturn Formation).	Middle Pennsylvanian...	Colorado.....
Huerfanito Bentonite Bed (of Lewis Shale).	Late Cretaceous.....	New Mexico.....
Hygiene Sandstone Member (of Pierre Shale).	Late Cretaceous.....	Colorado.....

Revision and reference

Friars Formation adopted as uppermost formation of La Jolla Group (not new). Rocks were formerly part of Hanna's (1926) Rose Canyon Shale Member of La Jolla Formation of Clark (1926). Overlies Scripps Formation (new). (Kennedy and Moore, 1971.)

Geographically extended into northwestern Georgia. (Cressler, 1970.)

Rocks of Van Buren Formation included in Gasconade Dolomite except in southeastern part of State. There, in its type locality, Van Buren remains in good usage. (Gann and others, 1971.)

Geertsen Canyon Quartzite adopted as uppermost formation in Brigham Group (newly revised); overlies Browns Hole Formation (new). Rocks formerly called Brigham Quartzite (restricted). (Crittenden and others, 1971.)

Gibson Jack Formation adopted. Overlies Camelback Mountain Formation (new). (Crittenden and others, 1971.)

Golden Door Volcanics is same rock unit as Patsy Mine Volcanics; Golden Door abandoned in favor of Patsy Mine Volcanics. (Anderson, 1971.)

Age changed from Silurian and Devonian to Middle and Late Silurian to Early Devonian. (Hatch and others, 1970.)

Age changed from early Tertiary(?) to Paleocene. (Phillips and others, 1971.)

Age changed from Devonian or older to Mississippian(?) or older. (Quinn, 1971.)

Graydon Channel Sandstone abandoned. (Gann and others, 1971.)

Greenfield Dolomite Member of Bass Islands Dolomite raised to formational rank in Bass Islands Group. (Norris and Fidler, 1971.)

Removed from Alder Group and made basal unit of Big Bug Group (new). Alder Group restricted to its type locality. (Anderson and others, 1971.)

Guaracanal Andesite redefined as Guaracanal Formation in northeastern Puerto Rico. Age changed from Paleocene and early Eocene(?) to Paleocene. (Seiders, 1971.)

In northeastern Puerto Rico, Martín Gonzáles Lava Member of Guaynabo Formation removed from Guaynabo and raised to formational rank. Leprocomio Siltstone Member removed from Guaynabo Formation and assigned to Frailes Formation as Leprocomio Mudstone Member. Guaynabo Formation not used in report. (Seiders, 1971.)

Gunsight Pass Member adopted; underlies and overlies unnamed upper unit and unnamed shale unit of Pierre, respectively. (Izett and others, 1971.)

In Missouri, Gunter Sandstone Member changed to member of Gasconade Dolomite except in southeastern part of State where it remains member of Van Buren Formation. Usage remains unchanged elsewhere. (Gann and others, 1971.)

Harkers Funglomerate of Slentz (1955) adopted and renamed Harkers Alluvium. (Tooker and Roberts, 1971a.)

Reduced in rank to member of Floyd Shale in northwestern Georgia. (Cressler, 1970.)

Hato Puerco Tuff redefined and name changed to Hato Puerco Formation. Age changed from Late Cretaceous(?) to Late Cretaceous. (Seiders, 1971.)

Age changed from Ordovician to Devonian to late Precambrian or Paleozoic. (Bryant and Reed, 1970.)

Age changed from Mississippian to Early Mississippian. (Sable, 1970.)

Age changed from Pliocene(?) to late Miocene and Pliocene(?). (Krieger and others, 1971.)

Age changed from late Miocene and Pliocene(?) to late Miocene and early Pliocene. (McKee and Anderson, 1971.)

Geographically extended into northwestern Georgia. (Cressler, 1970.)

Age changed from Pennsylvanian to Middle Pennsylvanian. (Mallory, 1971.)

Huerfanito Bentonite Bed adopted. (Fassett and Hinds, 1971.)

Hygiene Sandstone Member extended into north-central Colorado; underlies and overlies two unnamed shale members of Pierre. (Izett and others, 1971.)

Name	Age	Location
Idaho Canyon Tuff.....	late Miocene.....	Northwestern Nevada and southeastern Oregon.
Imuruk Volcanics.....	late Tertiary to middle Pleistocene.	Alaska.....
Infierno Formation.....	Late Cretaceous.....	Northeastern Puerto Rico.
Inkom Formation.....	late Precambrian.....	Southeastern Idaho (Pocatello area) and possibly northern Utah (Huntsville area).
Iron King Volcanics (of Alder Group).	Precambrian.....	Central Arizona.....
Jacque Mountain Limestone Member (of Minturn Formation).	Middle Pennsylvanian...	Colorado.....
Juana Diaz Formation.....	early and middle Oligocene.	Puerto Rico.....
Keeler Canyon Formation.....	Middle Pennsylvanian to Early Permian.	California.....
Kelley Canyon Formation.....	late Precambrian.....	Northern Utah (Huntsville area).
Kenilworth Member (of Blackhawk Formation).	Late Cretaceous.....	Utah.....
Kessler Limestone Member (of Bloyd Formation).	Early Pennsylvanian (Morrow).	Northeastern Oklahoma.
Kidder Limestone Member (of Monteagle Limestone).	Late Mississippian...	Kentucky and Tennessee.
Kirtland Shale.....	Late Cretaceous and Paleocene (in Mesa Portales area).	Northwestern New Mexico.
Kremmling Sandstone Member (of Pierre Shale).	Late Cretaceous...	North-central Colorado.
Krugloi Formation.....	Mesozoic or Tertiary....	Southwestern Alaska.
Kugruk Volcanics.....	late Tertiary and early Pleistocene(?).	Alaska.....
Kulthieth Formation.....	Paleocene to Oligocene..	Southern Alaska.....
La Jolla Group.....	early(?), middle, and late Eocene.	Southern California..
Latonia Shale.....	Late Ordovician.....	Kentucky and Ohio..
Lavender Shale Member (of Fort Payne Chert).	Early Mississippian (Osage).	Northwestern Georgia.
Leitchfield Formation.....	Late Mississippian (Chester).	Kentucky.....
Lenoir Limestone.....	Middle Ordovician.....	Northwestern Georgia.
Lenox Hills Formation.....	Early Permian (Wolfcamp).	Texas.....
Leprocomio Siltstone Member (of Guaynabo Formation).	Late Cretaceous.....	Northeastern Puerto Rico.
Lewis Shale.....	Late Cretaceous.....	New Mexico.....

Revision and reference

Idaho Canyon Tuff adopted. (Noble and others, 1970.)

Kugruk Volcanics abandoned; its rocks now included in Imuruk Volcanics. Age changed from early(?) and middle Pleistocene to late Tertiary to middle Pleistocene. (Hopkins and others, 1971.)

Infierno Formation adopted. Overlies Hato Puercio Formation; underlies Celada Formation (new). (Seiders, 1971.)

Inkom Formation adopted. In Huntsville area of northern Utah only, tentatively assigned to Brigham Group (newly revised). Overlies Caddy Canyon Quartzite (new) (only tentatively in Huntsville area); underlies Mutual Formation. (Crittenden and others, 1971.)

Removed from Alder Group, which is restricted to its type locality, and reassigned to the Big Bug Group (new). (Anderson and others, 1971.)

Age changed from Pennsylvanian to Middle Pennsylvanian. (Mallory, 1971.)

Age changed from middle Oligocene to early and middle Oligocene. (Glover, 1971.)

Age changed from Pennsylvanian and Early Permian to Middle Pennsylvanian to Early Permian. (Hall, 1971.)

Kelley Canyon Formation adopted. Overlies Maple Canyon Formation (new); underlies Caddy Canyon(?) Quartzite (new) (only tentatively in Huntsville area), lowermost formation of Brigham Group (newly revised). (Crittenden and others, 1971.)

Kenilworth Member of Young (1955) of Blackhawk Formation adopted. Overlies Aberdeen Member; underlies unnamed lower mudstone member. (Maberry, 1971.)

Geographically extended into northeastern Oklahoma. (Haley and Hendricks, 1971.)

Kidder Limestone Member adopted as upper member of Monteagle. Overlies Ste. Genevieve Limestone Member. (Lewis, 1971.)

In Mesa Portales area, age changed from Late Cretaceous to Late Cretaceous and Paleocene. (Fassett and Hinds, 1971.)

Kremmling Sandstone Member adopted. Underlies unnamed shale member of Pierre; overlies Sharon Springs Member of Pierre. (Izett and others, 1971.)

Krugloi Formation adopted. (Gates and others, 1971.)

Kugruk Volcanics abandoned; its rocks now included in overlying Imuruk Volcanics (revised). (Hopkins and others, 1971.)

Age changed from Eocene and Oligocene to Paleocene to Oligocene. (Addicott and Plaffer, 1971.)

La Jolla Formation of Clark (1926) adopted and raised in rank to La Jolla Group; includes (ascending): Mount Soledad Formation (new), Delmar Formation (not new), Torrey Sandstone (not new), Ardath Shale (new), Scripps Formation (new), and Friars Formation (new). (Kennedy and Moore, 1971.)

Latonia Shale abandoned; its rocks now included in Kope Formation. (Luft, 1971.)

Lavender Shale Member of Butts and Gildersleeve (1948) of Fort Payne Chert adopted. (Cressler, 1970.)

Vienna Limestone reduced in rank to member of Leitchfield Formation in Nolin Reservoir area. Remains in formational rank elsewhere. (Gildersleeve, 1971.)

Lenoir Limestone and its Mosheim Member geographically extended into northwestern Georgia. Deaton Formation of Spencer (1893) adopted as uppermost member of Lenoir in northwestern Georgia. (Cressler, 1970.)

Lenox Hills Formation of Ross (1959) adopted. Replaces, in part, Wolfcamp Formation, which is revised, in its type locality, to Wolfcamp Series. (Grant, 1971.)

Leprocomio Siltstone Member removed from Guaynabo Formation and reassigned to Frailes Formation as Leprocomio Mudstone Member in northeastern Puerto Rico. (Seiders, 1971.)

Huerfano Bentonite Bed (new) adopted and included in Lewis Shale. (Fassett and Hinds, 1971.)

Name	Age	Location
Lisburne Group.....	Mississippian, Pennsylvanian, and Permian.	Northern Alaska.....
Lomas Formation.....	Late Cretaceous.....	Northeastern Puerto Rico.
Lone Mountain Dolomite.....	Silurian (in Eureka area)	Nevada (Eureka area).
Lusardi Formation (of Rosario Group).	Late Cretaceous.....	Southern California..
McKinney Basalt (of Snake River Group).	Pleistocene.....	Southern Idaho.....
Magothy Formation.....	Late Cretaceous.....	New Jersey.....
Mahlac Member (of Alutom Formation).	early Miocene.....	Guam.....
Malpais Basalt.....	Pliocene.....	Nevada.....
Mancos Shale.....	Late Cretaceous.....	New Mexico.....
Maple Canyon Formation.....	late Precambrian.....	Northern Utah (Huntsville area).
Marcellus Shale.....	Middle Devonian.....	Pennsylvania and New Jersey.
Martín González Lava Member (of Guaynabo Formation).	Late Cretaceous.....	Northeastern Puerto Rico.
Massacre Bay Formation.....	late Tertiary or early Pleistocene.	Southwestern Alaska..
Maury Formation.....	Early Mississippian.....	Northwestern Georgia.
Middle Canyon Formation.....	Late Mississippian.....	South-central Idaho..
Middlefield Granite.....	Early(?) Devonian.....	Massachusetts.....
Milton Formation.....	Jurassic.....	California.....
Minturn Formation.....	Middle Pennsylvanian..	Colorado.....
Mission Valley Formation (of Poway Group).	late Eocene.....	Southern California..
Mitchell Creek Formation.....	Pennsylvanian.....	Northern Nevada.....
Moberly Channel Sandstone.....	Pennsylvanian.....	Missouri.....
Monacillo Formation.....	Late Cretaceous and early Tertiary(?).	Northeastern Puerto Rico.
Monk Formation.....	Precambrian.....	Northeastern Washington.
Monotony Tuff.....	late Oligocene.....	Nevada.....
Monteagle Limestone.....	Late Mississippian.....	Kentucky and Tennessee.
Moreno Formation.....	Late Cretaceous and Paleocene.	California.....
Mosheim Member (of Lenoir Limestone).	Middle Ordovician.....	Northwestern Georgia.
Mount Roberts Formation....	Pennsylvanian(?).....	Northwestern Washington.
Mount Soledad Formation (of La Jolla Group).	early(?) and middle Eocene.	Southern California..
Mount Stuart Granodiorite....	Late Cretaceous.....	Central Washington..
Muddy Buttes Sandstone Member (of Pierre Shale).	Late Cretaceous.....	North-central Colorado.

Revision and reference

Age changed from Early Mississippian to Middle Pennsylvanian to Mississippian, Pennsylvanian, and Permian. (Brosgé and TAILLEUR, 1971.)
Lomas Formation adopted; restricted to fault blocks. (Seiders, 1971.)

Age changed from Silurian and Early Devonian(?) (locally) to Silurian in Eureka area. (Nolan and others, 1971.)

Lusardi Formation of Nordstrom (1970) adopted as basal formation of Rosario Group; underlies Point Loma Formation (new). (Kennedy and Moore, 1971.)

Age changed from Holocene to Pleistocene. (Malde, 1971.)

Old Bridge Sand Member removed from Raritan Formation and made basal member of Magothy Formation. (Wolfe and Pakiser, 1971.)

Age changed from Oligocene to early Miocene. (Bukry and others, 1971.)

Age changed from Pliocene or Pleistocene to Pliocene. (Hedge and Noble, 1971.)

Whitewater Arroyo Shale Member of Owen (1966) of Dakota Sandstone adopted as Whitewater Arroyo Tongue of Mancos Shale. Twowells Sandstone Lentil of Pike (1947) of Mancos Shale adopted as Twowells Tongue of Dakota Sandstone. (Green and Pierson, 1971.)

Maple Canyon Formation adopted. Underlies Kelley Canyon Formation (new). (Crittenden and others, 1971.)

Brodhead Creek Member of Willard (1938) and Stony Hollow Member of Cooper (1941) adopted as upper and middle members, respectively, of Marcellus Shale in report area. Stony Hollow overlies Union Springs Shale Member, basal member of Marcellus. (Alvord and Drake, 1971.)

In northeastern Puerto Rico, Martín González Lava Member raised to formational rank as Martín González Lava; Guaynabo Formation not used. Usage remains unchanged elsewhere. (Seiders, 1971.)

Massacre Bay Formation adopted. (Gates and others, 1971.)

Reduced in rank to member of Chattanooga Shale in northwestern Georgia. (Cressler, 1970.)

Middle Canyon Formation of Huh (1967) adopted. (Mamet and others, 1971.)

Age changed from late Carboniferous or post-Carboniferous to Early(?) Devonian. (Hatch and others, 1970.)

Age changed from Triassic and Jurassic to Jurassic. (James, 1971.)

Age changed from Pennsylvanian (Des Moines and younger) to Middle Pennsylvanian. (Mallory, 1971.)

Mission Valley Formation adopted as middle formation of Poway Group. Overlies Stadium Conglomerate (new); underlies unnamed formation. (Kennedy and Moore, 1971.)

Mitchell Creek Formation adopted. (Coats, 1971.)

Moberly Channel Sandstone abandoned. (Gann and others, 1971.)

Age changed from Late Cretaceous to Late Cretaceous and early Tertiary(?). (Seiders, 1971.)

Age changed from Cambrian(?) to Precambrian. (Yates, 1970.)

Monotony Tuff adopted. (Ekren and others, 1971.)

Kidder Limestone adopted as upper member of Montegale Limestone, overlying Ste. Genevieve Limestone Member. (Lewis, 1971.)

Age changed from Late Cretaceous and Paleocene(?) to Late Cretaceous and Paleocene. (Bukry and others, 1971.)

Geographically extended into northwestern Georgia. (Cressler, 1970.)

Mount Roberts Formation of LeRoy (1913) adopted as used by Little (1960). (Yates, 1971.)

Mount Soledad Formation adopted as basal formation of La Jolla Group (not new); underlies Delmar Formation (not new). (Kennedy and Moore, 1971.)

Age changed from pre-Tertiary to Late Cretaceous. (Engels and Crowder, 1971.)

Muddy Buttes Sandstone Member adopted; underlies and overlies two unnamed shale members of Pierre. (Izett and others, 1971.)

Name	Age	Location
Mutual Formation.....	late Precambrian.....	Northern Utah (Huntsville area) and southeastern Idaho (Pocatello area).
Nancy Member (of Borden For- mation).	Early Mississippian.....	Kentucky.....
Neal Ranch Formation.....	Early Permian (Wolf- camp).	Texas.....
Nevidiskov Formation.....	middle Tertiary.....	Southwestern Alaska..
New Pass Tuff.....	Miocene.....	Central Nevada.....
New Providence Shale.....	Early Mississippian.....	Kentucky.....
Niobrara Formation.....	Late Cretaceous.....	Northwestern Colorado.
Northbridge Granite Gneiss....	early Paleozoic.....	Rhode Island and Massachusetts.
Norwood Tuff.....	Eocene and Oligocene....	Utah.....
Old Bridge Sand Member (of Raritan Formation).	Late Cretaceous.....	New Jersey.....
Ottawanah Rhyolite.....	Eocene or Oligocene....	Northern Nevada.....
Ox Frame Volcanics.....	Triassic.....	Southern Arizona....
Papoose Creek Formation.....	late Precambrian.....	Southeastern Idaho (Pocatello area).
Patapsco Formation.....	Early and Late Creta- ceous.	Maryland, Delaware, Virginia, Pennsyl- vania, and New Jersey.
Peabody Granite.....	Late Ordovician.....	Massachusetts.....
Peoria Loess.....	late Pleistocene.....	Western Kentucky....
Perkinsville Formation.....	Pliocene or Pleistocene..	Central Arizona.....
Perkinsville Formation.....	late Pliocene.....	Central and north- central Arizona.
Perryville Limestone Member (of Lexington Limestone).	Middle Ordovician.....	Kentucky.....
Pierre Shale.....	Late Cretaceous.....	North-central Colorado.
Pinto Diorite.....	Precambrian.....	Montana.....
Pinyon Conglomerate.....	Late Cretaceous and Paleocene.	Northwestern Wyo- ming (Yellowstone Park area).
Piper Formation.....	Middle Jurassic.....	Central Montana.....
Pleasanton Formation.....	Late Pennsylvanian (Missouri).	Missouri.....
Pocatello Formation.....	late Precambrian.....	Southeastern Idaho (Pocatello area).
Point Loma Formation (of Ro- sario Group).	Late Cretaceous.....	Southern California..
Point Pleasant Formation.....	Middle and Late Ordovician.	Northernmost Kentucky.

Revision and reference

- Geographically extended to Pocatello area of southeastern Idaho. Overlies Inkomo Formation (new); underlies Camelback Mountain Formation (new). In Huntsville area of northern Utah only, Mutual assigned to Brigham Group (newly revised), overlying Inkomo(?) Formation and underlying Browns Hole Formation (new) of Brigham Group. (Crittenden and others, 1971.)
- New Providence Shale reinstated in central Kentucky as basal member of Borden Formation. Nancy Member restricted to more silty, nonplastic shale within Nancy Member of previous usage. (Taylor and Lewis, 1971.)
- Neal Ranch Formation of Ross (1959) adopted. Replaces, in part, Wolfcamp Formation, which is revised, in its type locality, to Wolfcamp Series. (Grant, 1971.)
- Nevidskov Formation adopted. (Gates and others, 1971.)
- New Pass Tuff adopted. (McKee and Stewart, 1971.)
- New Providence Shale reinstated in central Kentucky as basal member of Borden Formation. Nancy Member restricted to more silty, nonplastic shale within Nancy Member of previous usage. (Taylor and Lewis, 1971.)
- Reduced in rank and included as member of Mancos Shale in northwestern Colorado. (Izett and others, 1971.)
- Age changed from Precambrian to early Paleozoic. (Quinn, 1971.)
- Age changed from Oligocene to Eocene and Oligocene. (Mullens, 1971.)
- Removed from Raritan Formation and included as basal member of Magothy Formation. (Wolfe and Pakiser, 1971.)
- Ottawanah Rhyolite adopted. (Coats, 1971.)
- Ox Frame Formation of Lootens (1965, 1966) adopted as Ox Frame Volcanics. Unconformably overlies Paleozoic rocks; disconformably underlies several Mesozoic formations. (Cooper, 1971.)
- Papoose Creek Formation adopted. Overlies Blackrock Canyon Limestone (new); underlies Caddy Canyon Quartzite (new). (Crittenden and others, 1971.)
- Geographically extended into New Jersey. Age changed from Early Cretaceous to Early and Late Cretaceous. (Wolfe and Pakiser, 1971.)
- Age changed from late(?) Paleozoic to Late Ordovician. (Zartman and Marvin, 1971.)
- Geographically extended into western Kentucky. (Finch, 1971.)
- Age changed from Pliocene(?) to Pleistocene(?) to Pliocene or Pleistocene. (Krieger and others, 1971.)
- Age changed from Pliocene or Pleistocene to late Pliocene. (McKee and Anderson, 1971.)
- Cornishville Member (of Perryville Formation) of Foerste (1912) adopted as Cornishville Bed of Perryville Limestone Member of Lexington Limestone. (Wolcott and Cressman, 1971.)
- In north-central Colorado, Pierre Shale includes (ascending): lower unit, Sharon Springs Member, Kremmling Sandstone Member (new), shale member, Muddy Buttes Sandstone Member (new), shale member, Hygiene Sandstone Member, shale member, Carter Sandstone Member (new), shale member, Gunsight Pass Member (new), and upper unit. (Izett and others, 1971.)
- Age changed from post-Cretaceous(?) to Precambrian. (Witkind, 1971.)
- Age changed from Paleocene to Late Cretaceous and Paleocene. Basal contact changed from bottom to top of conglomerate. (McKenna and Love, 1970.)
- Name changed to Piper Limestone in central Montana. (Witkind, 1971.)
- Warrensburg Channel Sandstone reduced in rank and assigned to Pleasanton Formation as Warrensburg Sandstone Member. (Gann and others, 1971.)
- Pocatello Formation of Ludlum (1942) adopted. Includes (ascending): unnamed lower member, Bannock Volcanic Member (not new), Scout Mountain Member (new), and unnamed upper member. Underlies Blackrock Canyon Limestone (new). (Crittenden and others, 1971.)
- Point Loma Formation adopted as middle formation of Rosario Group. Overlies Lusardi Formation (not new); underlies Cabrillo Formation (new). (Kennedy and Moore, 1971.)
- Bromley Shale of Bassler (1906) adopted as Bromley Shale Member of Point Pleasant Formation. (Luft, 1971.)

Name	Age	Location
Potomac Group	Early and Late Cretaceous.	Maryland, New Jersey, Virginia, Delaware, and Pennsylvania.
Poway Conglomerate	middle(?) and late Eocene.	Southern California..
Put-in-Bay Dolomite Member (of Bass Islands Dolomite.)	Late Silurian.....	Northern Ohio and southeastern Michigan.
Quail Porphyry	early Tertiary	Colorado
Quincy Granite	Late Ordovician	Massachusetts and Rhode Island.
Quinnville Quartzite (of Blackstone Series).	Precambrian(?)	Rhode Island
Quivira Shale Member (of Cherryvale Formation).	Late Pennsylvanian.....	Western Missouri.....
Raisin River Dolomite Member (of Bass Islands Dolomite).	Late Silurian.....	Northern Ohio and southeastern Michigan.
Raritan Formation	Late Cretaceous.....	Maryland and New Jersey.
Red Boy Rhyolite	Late Cretaceous	Southern Arizona.....
Resolution Dolomite Member (of Minturn Formation).	Middle Pennsylvanian...	Colorado
Robinson Limestone Member (of Minturn Formation).	Middle Pennsylvanian...	Colorado
Rockmart Slate	Middle Ordovician.....	Northwestern Georgia.
Rodolfo Formation	Triassic	Southern Arizona.....
Rosario Group	Late Cretaceous	Southern California..
Round Pass Mudflow	Holocene	Washington.....
Rowe Schist	Early to Late Cambrian and Early Ordovician.	Massachusetts, Vermont, and New York.
Roxana Silt	late Pleistocene	Western Kentucky...
Sabana Hoyos Limestone Member (of Cariblanco Formation).	Late Cretaceous	Puerto Rico
San Diego Lapilli Tuff Member (of Coamo Formation).	Late Cretaceous.....	Puerto Rico
Sandpoint Conglomerate	Late(?) Cretaceous	Idaho
Santa Ana Limestone Member (of Coamo Formation).	Late Cretaceous.....	Puerto Rico
Scott Peak Formation	Late Mississippian.....	Idaho
Scout Mountain Member (of Pocatello Formation).	late Precambrian.....	Southeastern Idaho (Pocatello area).
Scripps Formation (of La Jolla Group).	middle and late Eocene..	Southern California..

Revision and reference

Age changed from Early Cretaceous to Early and Late Cretaceous. (Wolfe and Pakiser, 1971.)

Redefinition of Poway Conglomerate by Hanna (1926) accepted; rank raised to Poway Group. Includes (ascending): Stadium Conglomerate (new), Mission Valley Formation (new), and unnamed formation. Age changed from late Tertiary to middle(?) and late Eocene. (Kennedy and Moore, 1971.)

Put-in-Bay Dolomite Member of Bass Islands Dolomite raised to formational rank in Bass Islands Group in Ohio; member rank is retained in Michigan. (Norris and Fidler, 1971.)

Age changed from Late Cretaceous or early Tertiary to early Tertiary. (Bergendahl and Koschmann, 1971.)

Age changed from Devonian(?) to Late Ordovician. (Zartman and Marvin, 1971.)

Quinnville Quartzite adopted. (Quinn, 1971.)

Geographically extended into western Missouri. (Connor and others, 1971.)

Raisin River Dolomite Member of Bass Islands Dolomite raised to formational rank in Bass Islands Group in Ohio; member rank is retained in Michigan. (Norris and Fidler, 1971.)

Excluded from Maryland; so-called Raritan in Maryland now considered upper part of Patapsco Formation. Old Bridge Sand Member removed from Raritan Formation in New Jersey and made basal member of Magothy Formation. (Wolfe and Pakiser, 1971.)

Red Boy Rhyolite of Thoms (1966, 1967) adopted. Mainly overlies Demetrie Volcanics (not new) and, to lesser extent, Ox Frame Volcanics (not new); is uppermost Cretaceous formation in Tascuela area. (Cooper, 1971.)

Age changed from Pennsylvanian to Middle Pennsylvanian. (Mallory, 1971.)

Age changed from Pennsylvanian to Middle Pennsylvanian. (Mallory, 1971.)

Rockmart Slate of Hayes (1891) adopted. (Cressler, 1970.)

Rodolfo Formation adopted. Unconformably overlies Paleozoic rocks; underlies Whitcomb Quartzite (new). (Cooper, 1971.)

Rosario Formation of Beal (1924) and Anderson and Hanna (1935) extended into southern California from Baja California and raised to group rank. Includes (ascending): Lusardi Formation (not new), Point Loma Formation (new), and Cabrillo Formation (new). (Kennedy and Moore, 1971.)

Round Pass Mudflow adopted. (Crandell, 1971.)

Age changed from Early to Late Cambrian and Early Ordovician(?) to Early to Late Cambrian and Early Ordovician. (Osberg and others, 1971.)

Geographically extended into western Kentucky. (Finch, 1971.)

Changed from member of Cariblanco Formation to member of Maravillas Formation. (Glover, 1971.)

Changed from San Diego Lapilli Tuff Member of Coamo Formation to San Diego Member of Maravillas Formation. (Glover, 1971.)

Sandpoint Conglomerate of Anderson (1930) adopted. (Harrison and Schmidt, 1971.)

Changed from member of Coamo Formation to member of Maravillas Formation. (Glover, 1971.)

Scott Peak Formation of Huh (1967) adopted. (Mamet and others, 1971.)

Scout Mountain Member adopted. Overlies Bannock Volcanic Member (not new); underlies unnamed upper member of Pocatello. (Crittenden and others, 1971.)

Scripps Formation of La Jolla Group (not new) adopted. Its rocks formerly part of Hanna's (1926) Rose Canyon Shale Member of La Jolla Formation of Clark (1926). Overlies Ardath Shale (new); underlies Friars Formation (new). (Kennedy and Moore, 1971.)

Name	Age	Location
Sehoo Formation.....	Pleistocene and Holocene	Nevada.....
Sharon Springs Member (of Pierre Shale).	Late Cretaceous.....	North-central Colorado.
Shingle Pass Tuff.....	early Miocene.....	Nevada.....
Shohola Member (of Catskill Formation).	Late Devonian.....	Northeastern Pennsylvania.
Silver Point Quartz Monzonite.	Tertiary.....	Northeastern Washington.
Sloane Peak Member (of State Bridge Formation).	Early Triassic.....	Colorado.....
Snake River Group.....	Pleistocene and Holocene	Southern Idaho.....
Soldier Meadow Tuff.....	late Miocene.....	Northwestern Nevada.
Sophie Mountain Formation...	Late Cretaceous(?).....	Northwestern Washington.
South Creek Formation.....	Late Mississippian (Chester).	Idaho.....
Stadium Conglomerate (of Poway Group).	middle(?) and late Eocene.	Southern California..
Stansbury Formation.....	Late Devonian.....	Northwestern Utah..
Starvation Flat Quartz Mon- zonite.	Mesozoic(?).....	Northeastern Washington.
State Bridge Formation.....	Permian and Early Triassic.	Colorado.....
Stevens Mountain Rhyolite....	Late Triassic or Early Jurassic.	Southern Arizona....
Stony Hollow Member (of Mar- cellus Shale).	Middle Devonian.....	Southeastern New York and north- eastern Pennsyl- sylvania.
Sullivan Buttes Latite.....	late Oligocene and early Miocene.	Central Arizona.....
Summit Lake Tuff.....	late Miocene.....	Northwestern Nevada.
Sunnyside Member (of Black- hawk Formation).	Late Cretaceous.....	Utah.....
Surrett Canyon Formation....	Late Mississippian.....	South-central Idaho..
Tabonuco Formation.....	Early Cretaceous.....	Northeastern Puerto Rico.
Tahkandit Limestone.....	Early Permian.....	East-central Alaska..
Talladega Slate.....	Precambrian(?) to Carboniferous(?).	Northwestern Georgia.
Tascuela Red Beds.....	Triassic.....	Southern Arizona....
Temblor Formation/Sandstone.	Oligocene and Miocene..	California.....

Revision and reference

Age changed from Pleistocene to Pleistocene and Holocene. (Birkeland and others, 1971.)

Geographically extended into north-central Colorado. Underlies Kremmling Sandstone Member (new) of Pierre; overlies unnamed lower unit of Pierre. (Izett and others, 1971.)

Age changed from Miocene to early Miocene. (Ekren and others, 1971.)

Shohola Formation of Willard (1936) adopted as Shohola Member, upper of three members of Catskill Formation. Overlies Anomink Red Shale Member (not new). (Alvord and Drake, 1971.)

Age changed from Mesozoic(?) to Tertiary. (Miller, 1971.)

Sloane Peak Member adopted. (Freeman, 1971.)

Stratigraphic succession revised. Formations are (ascending): Madson Basalt, Sugar Bowl Gravel, Thousand Springs Basalt, Crownsnest Gravel, Sand Springs Basalt, Wendell Grade Basalt, McKinney Basalt, Melon Gravel, and, locally, Holocene lava flows. Bancroft Springs Basalt abandoned; its rocks now part of McKinney Basalt. In southeastern Idaho the group also includes Big Hole Basalt. (Malde, 1971.)

Soldier Meadow Tuff adopted. (Noble and others, 1970.)

Sophie Mountain Conglomerate of Bruce (1917) adopted as used by Little (1960) and renamed Sophie Mountain Formation. (Yates, 1971.)

South Creek Formation of Huh (1967) adopted. (Mamet and others, 1971.)

Stadium Conglomerate adopted as basal formation of Poway Group; underlies Mission Valley Formation (new). (Kennedy and Moore, 1971.)

Stansbury Formation of Stokes and Arnold (1958) adopted. Overlies Laketown Dolomite; underlies Gardner Formation. (Tooker and Roberts, 1971b.)

Starvation Flat Quartz Monzonite adopted. (Clark and Miller, 1968.)

Sloane Peak of Early Triassic age adopted as member of State Bridge Formation. State Bridge divided into (ascending): Permian sandstone of the Fryingpan River and unnamed beds; Lower Triassic Sloane Peak Member (new), unnamed beds, and coarse unit at Toner Creek. Permian South Canyon Creek Member recognizable at north edge of quadrangle and pinches out 2 miles south of north edge. (Freeman, 1971.)

Stevens Mountain Rhyolite of Thoms (1966, 1967) adopted. Overlies Tascuela Red Beds (not new); underlies Demetrie Volcanics (not new). (Cooper, 1971.)

Stony Hollow Member of Cooper (1941) adopted as middle member of Marcellus Shale. Overlies Union Springs Shale Member; underlies Brodhead Creek Member (not new). (Alvord and Drake, 1971.)

Sullivan Buttes Latite adopted. (Krieger and others, 1971.)

Summit Lake Tuff adopted. (Noble and others, 1970.)

Sunnyside Member of Young (1955) of Blackhawk Formation adopted. Overlies unnamed lower mudstone member; underlies unnamed upper mudstone member. (Maberry, 1971.)

Surrett Canyon Formation of Huh (1967) adopted. (Mamet and others, 1971.)

Tabonuco Formation adopted. Overlies unnamed sequence of volcanoclastic rocks; underlies Hato Puerco Formation. (Seiders, 1971.)

Age changed from Permian to Early Permian. (Brabb and Grant, 1971.)

Cheaha Sandstone Member of Talladega Slate geographically extended into north-western Georgia. (Cressler, 1970.)

Tascuela Red Beds of Thoms (1966, 1967) adopted. Overlies Ox Frame Volcanics (not new); underlies Stevens Mountain Rhyolite (not new) or Demetrie Volcanics (not new). (Cooper, 1971.)

Big Blue Serpentinous Member removed from uppermost part of Temblor Formation and redefined as Big Blue Formation, overlying now stratigraphically restricted Temblor Formation in New Idria area. (Lockwood, 1971.)

Name	Age	Location
Texas Gulch Formation.....	Precambrian.....	Central Arizona.....
Toma de Agua Vitrophyre Member (of Cambalache Formation).	Late Cretaceous.....	Northeastern Puerto Rico.
Torrey Sandstone (of La Jolla Group).	middle Eocene.....	Southern California..
Trujillo Alto Limestone Member (of Monacillo Formation).	Late Cretaceous and early Tertiary(?).	Northeastern Puerto Rico.
Turupah Formation.....	Holocene.....	Nevada.....
Twowells Tongue (of Dakota Sandstone).	Late Cretaceous.....	New Mexico.....
Twowells Tongue (of Dakota Sandstone).	Late Cretaceous.....	New Mexico.....
Tymochtee Formation.....	Late Silurian.....	Northern Ohio.....
Valley Springs Formation.....	Miocene.....	Eastern California.....
Van Buren Formation.....	Early Ordovician.....	Missouri.....
Van Duzer Limestone.....	Devonian(?).....	Northern Nevada.....
Verde Formation.....	late Pliocene.....	Central and north-central Arizona.
Vienna Limestone.....	Late Mississippian (Chester).	Kentucky.....
Wahoo Limestone.....	Early Pennsylvanian to Permian.	Northern Alaska.....
Warrensburg Channel Sandstone.	Pennsylvanian.....	Missouri.....
Wendell Grade Basalt (of Snake River Group).	Pleistocene.....	Southern Idaho.....
Westerville Limestone Member (of Cherryvale Formation).	Late Pennsylvanian.....	Kansas, Nebraska, Iowa, and Missouri.
Whitcomb Quartzite.....	Early Cretaceous(?).....	Southern Arizona.....
White Quail Limestone Member (of Minturn Formation).	Pennsylvanian.....	Colorado.....
White Quail Limestone Member (of Minturn Formation).	Middle Pennsylvanian...	Colorado.....
Whitewater Arroyo Tongue (of Mancos Shale).	Late Cretaceous.....	New Mexico.....
Whitewater Arroyo Tongue (of Mancos Shale).	Late Cretaceous.....	New Mexico.....
Witnet Formation.....	early Tertiary.....	California.....
Wolf Porphyry.....	Eocene(?).....	Montana.....
Wolfcamp Formation.....	Early Permian (Wolfcamp).	Texas.....
Wrangell Lava.....	Miocene to Holocene....	Southern Alaska.....
Yorktown Formation.....	late Miocene and early Pliocene.	Virginia, North Carolina, Maryland, South Carolina, and District of Columbia.

Revision and reference

Removed from Alder Group, which is restricted to its type locality. Overlies Big Bug Group (new). (Anderson and others, 1971.)

Toma de Agua Vitrophyre Member adopted with restricted occurrence at top of Cambalache Formation (new). (Seiders, 1971.)

Torrey Sand of Hanna (1926), member of La Jolla Formation of Clark (1926), adopted as Torrey Sandstone of La Jolla Group. Overlies Delmar Formation (not new); underlies Ardath Shale (new). (Kennedy and Moore, 1971.)

Age changed from Late Cretaceous to Late Cretaceous and early Tertiary(?). (Seiders, 1971.)

Age changed from Pleistocene to Holocene. (Birkeland and others, 1971.)

Twowells Sandstone Lentil of Pike (1947) of Mancos Shale adopted as Twowells Tongue of Dakota Sandstone. (Green and Pierson, 1971.)

Name changed from Twowells Tongue to Twowells Sandstone Tongue. (Dane and others, 1971.)

Assigned to Bass Islands Group. (Norris and Fidler, 1971.)

Age changed from Miocene(?) to Miocene. (Clark, 1970.)

In Missouri, Van Buren Formation is restricted to its type locality in southeastern part of State. Elsewhere in Missouri, these rocks are included in Gasconade Dolomite. Van Buren remains in good usage in Oklahoma, Kansas, and Arkansas. (Gann and others, 1971.)

Van Duzer Limestone of Decker (1962) adopted. (Coats, 1971.)

Age changed from Pliocene or Pleistocene to late Pliocene. (McKee and Anderson, 1971.)

Included as member of Leitchfield Formation in Nolin Reservoir area. Remains formation or member of Buffalo Wallow Formation elsewhere. (Gildersleeve, 1971.)

Age changed from Early and Middle Pennsylvanian to Early Pennsylvanian to Permian. (Brosge and Tailleir, 1971.)

Reduced in rank and name changed to Warrensburg Sandstone Member of Pleasanton Formation. (Gann and others, 1971.)

Age changed from Holocene to Pleistocene. (Malde, 1971.)

Geographically extended into western Missouri. (Connor and others, 1971.)

Whitcomb Quartzite adopted. Overlies Rodolfo Formation (new); underlies Angelica Arkose (new). (Cooper, 1971.)

Age changed from Pennsylvanian and Permian(?) to Pennsylvanian. (Bergendahl and Koschmann, 1971.)

Age changed from Pennsylvanian to Middle Pennsylvanian. (Mallory, 1971.)

Whitewater Arroyo Shale Member of Owen (1966) of Dakota Sandstone adopted as Whitewater Arroyo Tongue of Mancos Shale. (Green and Pierson, 1971.)

Name changed from Whitewater Arroyo Tongue to Whitewater Arroyo Shale Tongue. (Dane and others, 1971.)

Age changed from pre-middle Miocene to early Tertiary. (Dibblee and Louke, 1970.)

Age changed from post-Cretaceous to Eocene(?). (Witkind, 1971.)

Wolfcamp Formation, in its type locality, revised to Wolfcamp Series. Wolfcamp is now used everywhere as a time-stratigraphic term (provincial series) rather than a rock-stratigraphic term (formation). Neal Ranch and Lenox Hills Formations (not new) adopted for rocks formerly within the Wolfcamp Formation. (Grant, 1971.)

Age changed from Tertiary to Holocene to Miocene to Holocene. (MacKevett, 1970.)

Age changed from Miocene to late Miocene and early Pliocene. (Hazel, 1971.)

REFERENCES

- Addicott, W. O., and Plafker, George, 1971, Paleocene mollusks from the Gulf of Alaska Tertiary province—A significant new occurrence on the North Pacific rim: U.S. Geol. Survey Prof. Paper 750-B, p. B48-B52.
- Adegoke, O. S., 1969, Stratigraphy and paleontology of the marine Neogene formations of the Coalinga region, California: California Univ. Pubs. Geol. Sci., v. 80, 269 p.
- Alvord, D. C., and Drake, A. A., Jr., 1971, Geologic map of the Bushkill quadrangle, Pennsylvania-New Jersey: U.S. Geol. Survey Geol. Quad. Map GQ-908.
- Anderson, A. L., 1928, Portland cement materials near Pocatello, Idaho: Idaho Bur. Mines and Geology, Pamph. 28, 15 p.
- 1930, Geology and ore deposits of the Clark Fork district, Idaho: Idaho Bur. Mines and Geology Bull. 12, 132 p.
- Anderson, C. A., Blacet, P. M., Silver, L. T., and Stern, T. W., 1971, Revision of Precambrian stratigraphy in the Prescott-Jerome area, Yavapai County, Arizona: U.S. Geol. Survey Bull. 1324-C, p. C1-C16.
- Anderson, F. M., and Hanna, G. D., 1935, Cretaceous geology of Lower California: California Acad. Sci. Proc., 4th ser., v. 23, no. 1, p. 1-34.
- Anderson, R. E., 1971, Thin skin distension in Tertiary rocks of southeastern Nevada: Geol. Soc. America Bull., v. 82, no. 1, p. 43-58.
- Bassler, R. S. 1906, A study of the James types of Ordovician and Silurian Bryozoa: U.S. Natl. Mus. Proc., v. 30, no. 1442, p. 1-66.
- Beal, C. H., 1924, Informe sobre la exploración geologica de la Baja California: Boletín Petroleo, v. 17, p. 421.
- Bergendahl, M. H., and Koschmann, A. H., 1971, Ore deposits of the Kokomo-Tenmile district, Colorado: U.S. Geol. Survey Prof. Paper 652, 53 p.
- Birkeland, P. W., Crandell, D. R., and Richmond, G. M., 1971, Status of correlation of Quaternary stratigraphic units in the western conterminous United States: Jour. Quaternary Research, v. 1, no. 2, p. 208-227.
- Brabb, E. E., Bukry, David, and Pierce, R. L., 1971, Eocene (Refugian) nannoplankton in the Church Creek Formation near Monterey, central California: U.S. Geol. Survey Prof. Paper 750-C, p. C44-C47.
- Brabb, E. E., and Grant, R. E., 1971, Stratigraphy and paleontology of the revised type section for the Tahkandit Limestone (Permian) in east-central Alaska: U.S. Geol. Survey Prof. Paper 703, 26 p.
- Brew, D. A., 1971, Mississippian stratigraphy of the Diamond Peak area, Eureka County, Nevada: U.S. Geol. Survey Prof. Paper 661, 84 p.
- Brosigé, W. P., and Tailleux, I. L., 1971, Northern Alaska petroleum province: Am. Assoc. Petroleum Geologists Mem. 15, p. 68-99.
- Bruce, E. L., 1917, Geology and ore deposits of Rossland: British Columbia Dept. Mines Bull. 4, 35 p.
- Bryant, Bruce, and Reed, J. C., Jr., 1970, Geology of the Grandfather Mountain window and vicinity, North Carolina and Tennessee: U.S. Geol. Survey Prof. Paper 615, 190 p.
- Bukry, David, Douglas, R. G., Kling, S. A., and Krasheninnikov, V., 1971, Planktonic micro-fossil biostratigraphy of the northwestern Pacific Ocean: Deep Sea Drilling Proj. Initial Repts., v. 6, p. 1253-1300.
- Butts, Charles, and Gildersleeve, Benjamin, 1948, Geology and mineral resources of the Paleozoic area in northwest Georgia: Georgia Geol. Survey Bull. 54, 176 p.
- Clark, B. L., 1926, The Domengine horizon, middle Eocene of California: California Univ., Dept. Geol. Sci. Bull., v. 16, no. 5, p. 99-118.

- Clark, L. D., 1970, Geology of the San Andreas 15-minute quadrangle, Calaveras County, California: California Div. Mines and Geology Bull. 195, 23 p.
- Clark, L. D., and Miller, F. K., 1968, Preliminary geologic map of the Chewelah Mountain quadrangle, Stevens County, Washington: Washington Div. Mines and Geology Geol. Map GM-5, 2 sheets.
- Coats, R. R., 1971, Geologic map of the Owyhee quadrangle, Nevada-Idaho: U.S. Geol. Survey Misc. Geol. Inv. Map I-665.
- Connor, J. J., Erdman, J. A., Sims, J. D., and Ebens, R. J., 1971, Roadside effects on trace element content of some rocks, soils, and plants of Missouri, in D. D. Hemphill, ed., Trace substances in environmental health: Missouri Univ. 4th Ann. Conf. on Trace Substances in Environmental Health, Proc., p. 26-34.
- Cooper, G. A., 1941, New Devonian stratigraphic units: Washington Acad. Sci. Jour., v. 31, no. 5, p. 179-181.
- Cooper, J. R., 1971, Mesozoic stratigraphy of the Sierrita Mountains, Pima County, Arizona: U.S. Geol. Survey Prof. Paper 658-D, p. D1-D42.
- Crandell, D. R., 1971, Postglacial lahars from Mount Rainier volcano, Washington: U.S. Geol. Survey Prof. Paper 677, 75 p.
- Cressler, C. W., 1970, Geology and ground-water resources of Floyd and Polk Counties, Georgia: Georgia Geol. Survey Inf. Circ. 39, 95 p.
- Crittenden, M. D., Jr., Schaeffer, F. E., Trimble, D. E., and Woodward, L. A., 1971, Nomenclature and correlation of some upper Precambrian and basal Cambrian sequences in western Utah and southeastern Idaho: Geol. Soc. America Bull., v. 82, no. 3, p. 581-602.
- Dane, C. H., Landis, E. R., and Cobban, W. A., 1971, The Twowells Sandstone Tongue of the Dakota Sandstone and the Tres Hermanos Sandstone as used by Herrick (1900), western New Mexico: U.S. Geol. Survey Prof. Paper 750-B, p. B17-B22.
- Decker, R. W., 1962, Geology of the Bull Run quadrangle, Elko County, Nevada: Nevada Bur. Mines Bull. 60, 65 p.
- Dibblee, T. W., Jr., and Chesterman, C. W., 1953, Geology of the Breckenridge Mountain quadrangle, California: California Div. Mines Bull. 168, 56 p.
- Dibblee, T. W., Jr., and Louke, G. P., 1970, Geologic map of the Tehachapi quadrangle, Kern County, California: U.S. Geol. Survey Misc. Geol. Inv. Map I-607.
- Dibblee, T. W., Jr., and Warne, A. H., 1970, Geologic map of the Cummings Mountain quadrangle, Kern County, California: U.S. Geol. Survey Misc. Geol. Inv. Map I-611.
- Dickinson, W. R., 1965, Tertiary stratigraphy of the Church Creek area, Monterey County, California, in Short contributions to California geology: California Div. Mines and Geology Spec. Rept. 86, p. 25-44.
- Ekren, E. B., Anderson, R. E., Rogers, C. L., and Noble, D. C., 1971, Geology of northern Nellis Air Force Base Bombing and Gunnery Range, Nye County, Nevada: U.S. Geol. Survey Prof. Paper 651, 91 p.
- Engels, J. C., and Crowder, D. F., 1971, Late Cretaceous fission-track and potassium-argon ages of the Mount Stuart Granodiorite and Beckler Peak stock, North Cascades, Washington: U.S. Geol. Survey Prof. Paper 750-D, p. D39-D43.
- Englund, K. J., and Windolph, J. F., Jr., 1971, Geology of the Carter Caves Sandstone (Mississippian) in northeastern Kentucky: U.S. Geol. Survey Prof. Paper 750-D, p. D99-D104.
- Fassett, J. E., and Hinds, J. S., 1971, Geology and fuel resources of the Fruitland Formation and Kirtland Shale of the San Juan Basin, New Mexico and Colorado: U.S. Geol. Survey Prof. Paper 676, 76 p.

- Finch, W. I., 1971, Geologic map of part of the Hickman quadrangle, Fulton County, Kentucky, and Mississippi County, Missouri: U.S. Geol. Survey Geol. Quad. Map GQ-874.
- Foerste, A. F., 1912, *Strophomena* and other fossils from Cincinnati and Mohawkian horizons, chiefly in Ohio, Indiana, and Kentucky: Denison Univ. Sci. Lab. Bull. 17, p. 17-172.
- Freeman, V. L., 1971, Stratigraphy of the State Bridge Formation in the Woody Creek quadrangle, Pitkin and Eagle Counties, Colorado: U.S. Geol. Survey Bull. 1324-F, p. F1-F17.
- Gann, E. E., Harvey, E. J., Jeffery, H. G., and Fuller, D. L., 1971, Water resources of northeastern Missouri: U.S. Geol. Survey Hydrol. Inv. Atlas HA-372, 4 sheets.
- Gates, Olcott, Powers, H. A., and Wilcox, R. E., 1971, Geology of the Near Islands, Alaska: U.S. Geol. Survey Bull. 1028-U, p. U709-U822.
- Gildersleeve, Benjamin, 1971, Geologic map of the Nolin Reservoir quadrangle, western Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-895.
- Glover, Lynn, III, 1971, Geology of the Coamo area, Puerto Rico, and its relation to the volcanic arc-trench association: U.S. Geol. Survey Prof. Paper 636, 102 p.
- Gordon, Mackenzie, Jr., 1971, Biostratigraphy and age of the Carboniferous formations, in Brew, D. A., Mississippian stratigraphy of the Diamond Peak area, Eureka County, Nevada: U.S. Geol. Survey Prof. Paper 661, p. 34-55.
- Grant, R. E., 1971, Brachiopods in the Permian reef environment of west Texas, in Yochelson, E. L., ed., Reef organisms through time—North American Paleont. Convention, Chicago 1969, Proc., pt. J: Lawrence, Kans., Allen Press, p. 1444-1481.
- Green, M. W., and Pierson, C. T., 1971, Geologic map of the Thoreau NE quadrangle, McKinley County, New Mexico: U.S. Geol. Survey Geol. Quad. Map GQ-954.
- Haley, B. R., and Hendricks, T. A., 1971, Geology of the Van Buren and Lavaca quadrangles, Arkansas-Oklahoma: U.S. Geol. Survey Prof. Paper 657-A, p. A1-A41.
- Hall, W. E., 1971, Geology of the Panamint Butte quadrangle, Inyo County, California: U.S. Geol. Survey Bull. 1299, 67 p.
- Hanna, M. A., 1926, Geology of the La Jolla quadrangle, California: California Univ., Dept. Geol. Sci. Bull., v. 16, no. 7, p. 187-246.
- Harrison, J. E., and Schmidt, P. W., 1971, Geologic map of the Elmira quadrangle, Bonner County, Idaho: U.S. Geol. Survey Geol. Quad GQ-953.
- Hatch, N. L., Jr., Norton, S. A., and Clark, R. G., Jr., 1970, Geologic map of the Chester quadrangle, Hampden and Hampshire Counties, Massachusetts: U.S. Geol. Survey Geol. Quad. Map GQ-858.
- Hayes, C. W., 1891, The overthrust faults of the southern Appalachians: Geol. Soc. America Bull., v. 2, p. 141-154.
- Hazel, J. E., 1971, Ostracode biostratigraphy of the Yorktown Formation (upper Miocene and lower Pliocene) of Virginia and North Carolina: U.S. Geol. Survey Prof. Paper 704, 13 p.
- Hedge, C. E., and Noble, D. C., 1971, Upper Cenozoic basalts with high $\text{Sr}^{87}/\text{Sr}^{86}$ and Sr/Rb ratios, southern Great Basin, western United States: Geol. Soc. America Bull., v. 82, no. 12, p. 3503-3509.
- Herrick, C. L., 1900, Report of a geological reconnaissance in western Socorro and Valencia Counties, New Mexico: Am. Geologist, v. 25, no. 6, p. 331-346.

- Hopkins, D. M., Matthews, J. V., Wolfe, J. A., and Silberman, M. L., 1971, A Pliocene flora and insect fauna from the Bering Strait region: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 9, no. 3, p. 211-231.
- Huh, O. K., 1967, The Mississippian System across the Wasatch line, east central Idaho, extreme southwestern Montana, in *Montana Geol. Soc., Centennial basin of southwest Montana—18th Annual Field Conference, August, 1967, Guidebook: [Billings, Mont.] Montana Geol. Soc.*, p. 31-62.
- Izett, G. A., Cobban, W. A., and Gill, J. R., 1971, The Pierre Shale near Kremmling, Colorado, and its correlation to the east and the west: *U.S. Geol. Survey Prof. Paper 684-A*, p. A1-A19.
- James, O. B., 1971, Origin and emplacement of the ultramafic rocks of the Emigrant Gap area, California: *Jour. Petrology*, v. 12, p. 523-560.
- Kennedy, M. P., and Moore, G. W., 1971, Stratigraphic relations of Upper Cretaceous and Eocene formations, San Diego coastal area, California: *Am. Assoc. Petroleum Geologists Bull.*, v. 55, no. 5, p. 709-722.
- Krieger, M. H., Creasey, S. C., and Marvin, R. F., 1971, Ages of some Tertiary andesitic and latitic volcanic rocks in the Prescott-Jerome area, north-central Arizona: *U.S. Geol. Survey Prof. Paper 750-B*, p. B157-B160.
- LeRoy, O. E., 1913, West Kootenay and Boundary districts: *Internat. Geol. Cong.*, 12th, Canada, Guidebook 9, p. 61-102.
- Lewis, R. Q., Sr., 1971, The Monteagle Limestone of south-central Kentucky: *U.S. Geol. Survey Bull.* 1324-E, p. E1-E10.
- Little, H. W., 1960, Nelson map-area, west half, British Columbia: *Canada Geol. Survey Mem.* 308, 205 p.
- Lockwood, J. P., 1971, Sedimentary and gravity-slide emplacement of serpentinite: *Geol. Soc. America Bull.*, v. 82, no. 4, p. 919-936.
- Lootens, D. J., 1965, Structure and petrography of the east side of the Sierrita Mountains, Pima County, Arizona: Tucson, Ariz., Arizona Univ. unpub. Ph. D. dissert., 227 p. [Abs. in *Dissert. Abs.*, v. 26, no. 4, p. 2135.]
- 1966, Geology and structural environment of the east side of the Sierrita Mountains, Pima County, Arizona: *Arizona Geol. Soc. Digest*, v. 8, p. 33-56.
- Ludlum, J. C., 1942, Pre-Cambrian formations at Pocatello, Idaho: *Jour. Geology*, v. 50, no. 1, p. 85-95.
- 1943, Structure and stratigraphy of part of the Bannock Range, Idaho: *Geol. Soc. America Bull.*, v. 54, no. 7, p. 973-986.
- Luft, S. J., 1971, Geologic map of part of the Covington quadrangle, northern Kentucky: *U.S. Geol. Survey Geol. Quad. Map GQ-955*.
- Maberry, J. O., 1971, Sedimentary features of the Blackhawk Formation (Cretaceous) in the Sunnyside district, Carbon County, Utah: *U.S. Geol. Survey Prof. Paper 688*, 44 p.
- McKee, E. H., and Anderson, C. A., 1971, Age and chemistry of Tertiary volcanic rocks in north-central Arizona and relation of the rocks to the Colorado Plateaus: *Geol. Soc. America Bull.*, v. 82, no. 10, p. 2767-2782.
- McKee, E. H., and Stewart, J. H., 1971, Stratigraphy and potassium-argon ages of some Tertiary tuffs from Lander and Churchill Counties, central Nevada: *U.S. Geol. Survey Bull.* 1311-B, p. B1-B28.
- McKenna, M. C., and Love, J. D., 1970, Local stratigraphic and tectonic significance of *Leptoceratops*, a Cretaceous dinosaur in the Pinyon Conglomerate, northwestern Wyoming: *U.S. Geol. Survey Prof. Paper 700-D*, p. D55-D61.
- MacKevett, E. M., Jr., 1970, Geology of the McCarthy B-4 quadrangle, Alaska: *U.S. Geol. Survey Bull.* 1333, 31 p.

- Malde, H. E., 1971, History of Snake River Canyon indicated by revised stratigraphy of Snake River Group near Hagerman and King Hill, Idaho: U.S. Geol. Survey Prof. Paper 644-F, p. F1-F21.
- Mallory, W. W., 1971, The Eagle Valley Evaporite, northwest Colorado—A regional synthesis: U.S. Geol. Survey Bull. 1311-E, p. E1-E37.
- Mamet, B. L., Skipp, Betty, Sando, W. J., and Mapel, W. J., 1971, Biostratigraphy of Upper Mississippian and associated Carboniferous rocks in south-central Idaho: Am. Assoc. Petroleum Geologists Bull., v. 55, no. 1, p. 20-33.
- Miller, F. K., 1971, The Newport fault and associated mylonites, northeastern Washington: U.S. Geol. Survey Prof. Paper 750-D, p. D77-D79.
- Monroe, W. H., 1971, Geologic map of the Manati quadrangle, Puerto Rico: U.S. Geol. Survey Misc. Geol. Inv. Map I-671.
- Mullens, T. E., 1971, Reconnaissance study of the Wasatch, Evanston, and Echo Canyon Formations in part of northern Utah: U.S. Geol. Survey Bull. 1311-D, p. D1-D31.
- Noble, D. C., McKee, E. H., Smith, J. G., and Korringa, M. K., 1970, Stratigraphy and geochronology of Miocene volcanic rocks in northwestern Nevada: U.S. Geol. Survey Prof. Paper 700-D, p. D23-D32.
- Nolan, T. B., Merriam, C. W., and Brew, D. A., 1971, Geologic map of the Eureka quadrangle, Eureka and White Pine Counties, Nevada: U.S. Geol. Survey Misc. Geol. Inv. Map I-612.
- Nordstrom, C. E., 1970, Lusardi Formation; A post-batholithic Cretaceous conglomerate north of San Diego, California: Geol. Soc. America Bull., v. 81, no. 2, p. 601-605.
- Norris, S. E., and Fidler, R. E., 1971, Availability of ground water from limestone and dolomite aquifers in northwest Ohio and its relation to geologic structure: U.S. Geol. Survey Prof. Paper 750-B, p. B229-B235.
- Osberg, P. H., Hatch, N. L., Jr., and Norton, S. A., 1971, Geologic map of the Plainfield quadrangle, Franklin, Hampshire, and Berkshire Counties, Massachusetts: U.S. Geol. Survey Geol. Quad. Map GQ-877.
- Owen, D. E., 1966, Nomenclature of Dakota Sandstone (Cretaceous) in San Juan basin, New Mexico and Colorado: Am. Assoc. Petroleum Geologists, v. 50, no. 5, p. 1023-1028.
- Pavrides, Louis, 1971, Geologic map of the Houlton quadrangle, Aroostook County, Maine: U.S. Geol. Survey Geol. Quad. Map GQ-920.
- Phillips, C. H., Cornwall, H. R., and Rubin, Meyer, 1971, A Holocene ore body of copper oxides and carbonates at Ray, Arizona: Econ. Geology, v. 66, no. 3, p. 495-498.
- Pike, W. S., Jr., 1947, Intertonguing marine and nonmarine Upper Cretaceous deposits of New Mexico, Arizona, and southwestern Colorado: Geol. Soc. America Mem. 24, 103 p.
- Quinn, A. W., 1971, Bedrock geology of Rhode Island: U.S. Geol. Survey Bull. 1295, 68 p.
- Ross, C. A., 1959, The Wolfcamp Series (Permian) and new species of fusulinids, Glass Mountains, Texas: Washington Acad. Sci. Jour., v. 49, no. 9, p. 299-316.
- Sable, El G., 1970, Source areas of Lower Mississippian red beds in eastern mid-continent: U.S. Geol. Survey Prof. Paper 700-D, p. D88-D91.
- Seiders, V. M., 1971, Cretaceous and lower Tertiary stratigraphy of the Gurabo and El Yunque quadrangles, Puerto Rico: U.S. Geol. Survey Bull. 1294-F, p. F1-F58.
- Slentz, L. W., 1955, Salt Lake group in lower Jordan Valley, Utah, in Utah Geol. Soc., Guidebook to the geology of Utah, no. 10, Tertiary and Quaternary

- geology of the eastern Bonneville basin: Salt Lake City, Utah Geol. and Mineralog. Survey, p. 23-36.
- Southwick, D. L., Reed, J. C., Jr., and Mixon, R. B., 1971, The Chopawamsic Formation—A new stratigraphic unit in the Piedmont of northeastern Virginia: U. S. Geol. Survey Bull. 1324-D, p. D1-D11.
- Spencer, J. W., 1893, The Paleozoic group; the geology of ten counties of north-western Georgia: Atlanta, Ga., Georgia Geol. Survey, 406 p.
- Stokes, W. L., and Arnold, D. E., 1958, Northern Stansbury Range and the Stansbury Formation, in Utah Geol. Soc., Guidebook to the geology of Utah, no. 13, Geology of the Stansbury Mountains, Tooele County, Utah: Salt Lake City, Utah Geol. and Mineralog. Survey, p. 135-149.
- Taylor, A. R., and Lewis, R. Q., Sr., 1971, Geologic map of the Yosemite quadrangle, Casey and Lincoln Counties, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-910.
- Thoms, J. A., 1966, The geology and ore deposits of the Tascuela area, Sierrita Mountains, Pima County, Arizona: Ann Arbor, Mich., Univ. Michigan unpub. Ph. D. dissert., 371 p.
- 1967, The geology and ore deposits of the Tascuela area, Sierrita Mountains, Pima County, Arizona [abs.]: Dissert. Abs., Sec. B, Sci. and Eng., v. 27, no. 7, p. 2420B-2421B.
- Tooker, E. W., and Roberts, R. J., 1971a, Geologic map of the Magna quadrangle, Salt Lake County, Utah: U.S. Geol. Survey Geol. Quad. Map GQ-923.
- 1971b, Structures related to thrust faults in the Stansbury Mountains, Utah: U.S. Geol. Survey Prof. Paper 750-B, p. B1-B12.
- Weir, G. W., Gualtieri, J. L., and Schlanger, S. O., 1966, Borden Formation (Mississippian) in south- and southeast-central Kentucky: U.S. Geol. Survey Bull. 1224-F, 38 p.
- Wells, J. D., Elliott, J. E., and Obradovich, J. D., 1971, Age of the igneous rocks associated with ore deposits, Cortez-Buckhorn area, Nevada: U.S. Geol. Survey Prof. Paper 750-C, p. C127-C135.
- White, I. C., 1882, The geology of Pike and Monroe Counties, Pennsylvania: Pennsylvania Geol. Survey, 2d ser., Rept. G6, 407 p.
- Willard, Bradford, 1935, Portage Group in Pennsylvania: Geol. Soc. America Bull., v. 46, no. 8, p. 1195-1218.
- 1936, Continental Upper Devonian of northeastern Pennsylvania: Geol. Soc. America Bull., v. 47, no. 4, p. 565-607.
- 1937, Devonian nomenclature in Pennsylvania: Pennsylvania Acad. Sci. Proc., v. 11, p. 26-34.
- 1938, A Paleozoic section at Delaware Water Gap: Pennsylvania Geol. Survey, 4th ser., Bull. G-11, 35 p.
- 1939, The Devonian of Pennsylvania: Middle and Upper Devonian: Pennsylvania Geol. Survey, 4th ser., Bull. G-19, p. 1-29, 131-481.
- Williams, N. C., and Madsen, J. H., Jr., 1959, Late Cretaceous stratigraphy of the Coalville area, Utah, in Intermountain Assoc. Petroleum Geologists, Guidebook to the geology of the Wasatch and Uinta Mountains transition area, 10th Annual Field Conference 1959: Salt Lake City, Utah, p. 122-125.
- Witkind, I. J., 1971, Geologic map of the Barker quadrangle, Judith Basin and Cascade Counties, Montana: U.S. Geol. Survey Geol. Quad. Map GQ-898.
- Wolcott, D. E., and Cressman, E. R., 1971, Geologic map of the Bryantsville quadrangle, central Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-945.

- Wolfe, J. A., and Pakiser, H. M., 1971, Stratigraphic interpretations of some Cretaceous microfossil floras of the Middle Atlantic States: U.S. Geol. Survey Prof. Paper 750-B, p. B35-B47.
- Yates, R. G., 1970, Geologic background of the Metaline and Northport mining districts, Washington, [Chap. 3] in *Lead-zinc deposits in the Kootenay Arc, northeastern Washington and adjacent British Columbia*—Soc. Econ. Geologists, 1970 Northwest Field Conf. Guidebook: Washington Div. Mines and Geology Bull. 61, p. 17-39.
- 1971, Geologic map of the Northport quadrangle, Washington: U.S. Geol. Survey Misc. Geol. Inv. Map I-603.
- Young, R. G., 1955, Sedimentary facies and intertonguing in the Upper Cretaceous of the Book Cliffs, Utah-Colorado: Geol. Soc. America Bull., v. 66, no. 2, p. 177-201.
- Zartman, R. E., and Marvin, R. F., 1971, Radiometric age (Late Ordovician) of the Quincy, Cape Ann, and Peabody Granites from eastern Massachusetts: Geol. Soc. America Bull., v. 82, no. 4, p. 937-958.
- Zen, E-an, and Ratcliffe, N. M., 1971, Bedrock geologic map of the Egremont quadrangle and adjacent areas, Berkshire County, Massachusetts, and Columbia County, New York: U.S. Geol. Survey Misc. Geol. Inv. Map I-628.